

nigris : loris et supercilio lato usque ad latera colli producto cinereis. Long. tot. 6·1 poll., culm. 0·7, alæ 3·0, caudæ 2·2, tarsi 1·1.

Hab. Efulen, Cameroons (*G. W. Bates*). Como River, 60 miles from Gaboon (*G. W. Bates*). R. Congo (*Petit : Mus. Brit.*).

Obs. A species easily distinguished from all the other African members of the genus by its black head and grey eyebrow.

2. TURDINUS CERVINIVENTRIS, n. sp.

Turdinus fulvescens, partim, auct. recent. (nec Cass.).

♂. Similis *T. fulvescenti*, Cass., sed corpore subtus cervino-brunneo, abdomine vix pallidiore : gula albida, griseo obscure striata : subcaudalibus cervinis, vix rufescentibus. Long. tot. 6·7 poll., culm. 0·8, alæ 2·95, caudæ 2·6, tarsi 0·95.

Hab. Conde, Congo (*Petit*), Chissambo (*Petit*), Yambuya, Aruwimi River (*Jameson*), Como River, Gaboon (*G. L. Bates*), Jaunde, Camaroons (*Zenker*).

Obs. The true *T. fulvescens*, Cass., of which the Museum contains two co-types, is a white-bellied species. Cassin appears to have confused two distinct species under the heading of *T. fulvescens*, as he observes : " In some specimens the entire underparts are dull reddish fulvous." These would be the new *T. cerviniventris*.

3. SCOPS HOLERYTHRA, n. sp.

Similis *S. icterorhynchæ*, Shelley, sed ubique cinnamomea, maculis albis scapularibus nullis, tectricibus alarum quoque albo haud maculatis : subtus læte cinnamomea, maculis longitudinalibus albis notata, sed minime fusco-vermiculata. Long. tot. 8·5 poll., culm. 0·8, alæ 5·5, caudæ 2·9, tarsi 1·05.

Hab. Efulen, Kamerun.

4. HAPALODERMA ÆQUATORIALE, n. sp.

Similis *H. narinæ*, sed minor : alæ fasciis vermiculatis nigris fortissime indicatis. Long. tot. 10·5 poll., culm. 0·8, alæ 4·7, caudæ 5·8, tarsi 0·65.

Hab. Efulen, Kamerun.