

Differs from *Ninox jacquinoti eichhorni* by the reduction of the whitish bars on the wing; wing, 190-196 mm.

RANGE.—Mono or Treasury Island, Solomon Islands.

***Tyto alba interposita*, new subspecies**

TYPE.—No. 215566, Amer. Mus. Nat. Hist.; ♂ ad.; Vanikoro, Santa Cruz Islands; September 21, 1926; R. H. Beck and J. G. Correia.

Differs from the neighboring races by the buffy to ochraceous wash of the under parts and the prevalence of the orange-ochre color throughout the plumage; wing, 268-279 mm.

RANGE.—Santa Cruz Islands, Banks Islands, and Northern New Hebrides.

***Tyto alba crassirostris*, new subspecies**

TYPE.—No. 335322, Amer. Mus. Nat. Hist.; ♀ ad.; Boang Island, Tanga group, Bismarck Archipelago; February 4, 1935, W. F. Coultas.

Differs from *T. a. delicatula* by the stronger bill and feet, and the darker coloration; wing, 285-290 mm.

RANGE.—Known only from the type-locality.

***Collocalia pelewensis*, new species**

TYPE.—No. 332022, Amer. Mus. Nat. Hist.; ♀ ad.; Palau Island, Micronesia; December 11, 1931; W. F. Coultas.

Small; tarsus naked; upper parts dark fuscous-green, with a brownish tone on the back; crown not very much darker than back; rump pale but no distinct light gray bar across the rump as in *spodiopygia*; there is much individual variation in the color of the rump, the bases of the feathers always being pale gray, the tips, however, sometimes being strongly glossy green, sometimes grayish; inner margins of the wing-feathers not particularly light; feathers of chin and throat soft and decomposed, with fuscous bases and rather sharply defined silvery gray edges, but no shaft-streaks; abdomen dull gray, slightly darker than throat, inconspicuous shaft-streaks on breast and abdomen, more pronounced ones on under tail-coverts; longest under tail-coverts fairly glossy green; white loreal spot inconspicuous. Wing, 109-115 mm.

RANGE.—Palau Islands.

***Collocalia vanikorensis ponapensis*, new subspecies**

TYPE.—No. 331343, Amer. Mus. Nat. Hist.; ♀ ad.; Ponape, Caroline Islands; November 22, 1930; W. F. Coultas.

Similar to *C. v. inquieta* (wing, 117-123 mm.) and *C. v. rukensis* (wing, 112-119.5 mm.), but smaller (wing, 106-113 mm.).

RANGE.—Ponape, Caroline Islands.

***Collocalia lowi orientalis*, new subspecies**

TYPE.—No. 225168, Amer. Mus. Nat. Hist.; ♂ ad.; Guadalcanar, Solomon Islands; May 28, 1927; R. H. Beck.