

- 392C. *TYTO NOVÆ-HOLLANDIÆ RIORDANI*, subsp. n.
Victorian White-faced Owl.

Differs from all other subspecies of *T. novæ-hollandiæ* in its darker upper-surface and larger-sized wing, 343 mm.

Type, Victoria (Warnambool), No. 11,104.

Range, Victoria.

- 394A. *TYTO NOVÆ-HOLLANDIÆ MELVILLENSIS*, subsp. n.
Melville Island Chestnut-faced Owl.

Differs from *T. n. perplexa* in its smaller size and darker buff below.

Type, Melville Island, Northern Territory, No. 10,795.

Range, Melville Island.

- 404A. *TRICHOGLOSSUS RUBRITORQUIS MELVILLENSIS*,
subsp. n.

Northern Red-collared Lorikeet.

Differs from *T. r. rubritorquis* in having the abdomen bluish instead of olive.

Type, Melville Island, Northern Territory, No. 10,876.

Range, Melville Island.

- 418A. *CALYPTORHYNCHUS FUNEREUS WHITEÆ*, subsp. n.
Kangaroo Island Black Cockatoo.

Differs from *C. f. funereus* in its smaller size: wing 281 (typical wing 314 mm.).

Type, Kangaroo Island, South Australia, No. 10,355.

Range, Kangaroo Island.

- 421A. *CALYPTORHYNCHUS BANKSII FITZROYI*, subsp. n.
Western Great-billed Cockatoo.

Differs from *C. b. macrorhynchus* in its much less massive bill.

Type, Fitzroy River, North-west Australia, No. 9,407.

Range, North-west Australia.